



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

APRIL 19.

J. CHESTON MORRIS, M. D., in the Chair.

Forty-seven persons present.

The Publication Committee reported the reception of papers under the following titles:

"*Stauroneis terryi* D. B. Ward," by T. Chalkley Palmer (April 11, 1910).

"A revision of the North American Species of the Genus *Ischnoptera* (Orthoptera)," by James A. G. Rehn and Morgan Hebard (April 12, 1910).

"A new Diatom," by T. Chalkley Palmer (April 19, 1910).

Under the auspices of the Biological and Microscopical Section the following communications were made:

Jelly-pores in the Diatomaceæ.—MR. CHARLES S. BOYER discussed the observations of Otto Müller, George Karsten and others upon the occurrence and function of the *jelly pore* in the Diatomaceæ and described it as found in certain species not heretofore recorded. As has been stated, the production of a stipe or jelly cushion appears to be its function. A more minute study of the habitat in various motile and immotile species is requisite. The speaker dwelt upon the varying conditions in earlier and later cell growth, of the tube and thallus forming jelly, and of the relation between the raphé and the pore. Suggestions were made as to the relation between the forms producing the stipes and jelly cushions and those in which the protective devices of many pelagic species were found to be siliceous. Drawings, original, and from various authors, illustrated the remarks.

DR. THOMAS S. STEWART, on the Hook-worm, *Ankylostoma duodenale*. (No abstract.)

Unusual Forms of Myxomycetes.—MR. HUGO BILGRAM remarked that in the early stages of their existence Myxomycetes are composed of a slimy mass consisting of cells resembling amœbæ, and only in the last hours of their life do they assume the apparently organized form of the well-known beautiful sporangia.

The most simple species consist of a mass of spores enclosed in a sporangium wall. Others develop within the sporangium a capillitium often consisting of branching filaments, and in some genera, of isolated fibres. In many genera the capillitium bears characteristic markings, such as spines, warts or spiral ribs. In the process of maturation